




Rabbit Anti-Human c-Met (Paired1003) Monoclonal Antibody (Clone SP59)

CATALOG #:	M3590	0.1 ml rabbit monoclonal antibody purified by protein A/G in PBS/1% BSA buffer pH 7.6 with less than 0.1% sodium azide.
	M3592	0.5 ml rabbit monoclonal antibody purified by protein A/G in PBS/1% BSA buffer pH 7.6 with less than 0.1% sodium azide.
	M3594	1.0 ml rabbit monoclonal antibody purified by protein A/G in PBS/1% BSA buffer pH 7.6 with less than 0.1% sodium azide.
	M3591	7 ml pre-diluted rabbit monoclonal antibody purified by protein A/G in TBS/1% BSA buffer pH 7.6 with less than 0.1% sodium azide.
INTENDED USE:	For Research Use Only. Not for use in diagnostic procedures.	
CLONE:	SP59	
IMMUNOGEN:	Synthetic peptide corresponding to the region of c-Met that contains non-phosphorylated tyrosine1003.	
IG ISOTYPE:	Rabbit IgG	
EPITOPE:	Not determined	
MOLECULAR WEIGHT	145 kDa	
SPECIES REACTIVITY:	Human (tested). (See www.springbio.com for information on species reactivity predicted by sequence homology.)	
DESCRIPTION:	The c-Met oncogene was originally isolated from chemical carcinogen-treated human osteogenic sarcoma cell line by transfection analysis in NIH/3T3 cells. The Met proto-oncogene product was identified as a transmembrane receptor-like protein with tyrosine kinase activity that is expressed in many tissues. The c-Met gene product has been identified as the cell surface receptor for hepatocyte growth factor, a plasminogen-like protein thought to be a humoral mediator of liver regeneration.	
APPLICATIONS:	Immunohistochemistry (IHC)	
IHC PROCEDURE:	<p>Specimen Preparation: Formalin-fixed, paraffin-embedded tissues are suitable for use with this primary antibody.</p> <p>Deparaffinization: Deparaffinize slides using xylene or xylene alternative and graded alcohols.</p> <p>Antibody Dilution: If using the concentrate format of this product, dilute the antibody 1:100. The dilutions are estimates; actual results may differ because of variability in methods and protocols.</p> <p>Antigen Retrieval: Boil tissue section in 10mM citrate buffer, pH 6.0 for 10 min followed by cooling at room temperature for 20 min.</p> <p>Primary Antibody Incubation: Incubate for 30 minutes at room temperature.</p> <p>Slide Washing: Slides must be washed in between steps. Rinse slides with PBS/0.05% Tween.</p> <p>Visualization: Detect the antibody as instructed by the instructions provided with the visualization system.</p>	
POSITIVE CONTROL:	Breast carcinoma, colon carcinoma	
CELLULAR LOCALIZATION:	Membrane	
STORAGE & STABILITY	<p>Store at 2-8°C. Do not freeze. The user must validate any other storage conditions. When properly stored, the reagent is stable to the date indicated on the label. Do not use the reagent beyond the expiration date.</p> <p>There are no definitive signs to indicate instability of this product; therefore, positive and negative controls should be tested simultaneously with unknown specimens.</p> <p>If unexpected results are observed which cannot be explained by variations in laboratory procedures and a problem with the reagent is suspected, contact Technical Support at spring.tech@ventana.roche.com.</p>	
WARNINGS & PRECAUTIONS:	<ol style="list-style-type: none"> 1. Avoid contact of reagents with eyes and mucous membranes. If reagents come into contact with sensitive areas, wash with copious amounts of water. 2. This product is harmful if swallowed. 3. Consult local or state authorities with regard to recommended method of disposal. 4. Avoid microbial contamination of reagents. 	



Human breast carcinoma stained with anti-c-Met (paired1003) antibody