




Rabbit Anti-Human *c-erbB2* Monoclonal Antibody (Clone SP3)

CATALOG #:	M3030	0.1 ml rabbit monoclonal antibody supplied as tissue culture supernatant in TBS/1% BSA buffer pH 7.5 with less than 0.1% sodium azide.
	M3032	0.5 ml rabbit monoclonal antibody supplied as tissue culture supernatant in TBS/1% BSA buffer pH 7.5 with less than 0.1% sodium azide.
	M3034	1.0 ml rabbit monoclonal antibody supplied as tissue culture supernatant in TBS/1% BSA buffer pH 7.5 with less than 0.1% sodium azide.
	M3031	7.0 ml pre-diluted rabbit monoclonal antibody supplied as tissue culture supernatant in TBS/1% BSA buffer pH 7.6 with less than 0.1% sodium azide.



Human breast carcinoma stained with anti-c-erbB2 antibody

INTENDED USE:	For Research Use Only. Not for use in diagnostic procedures.
CLONE:	SP3
IMMUNOGEN:	Recombinant protein encoding extracellular domain of human <i>c-erbB2</i> .
IG ISOTYPE:	Rabbit IgG
EPITOPE:	Not determined
MOLECULAR WEIGHT	185kDa
SPECIES REACTIVITY:	Human (tested). (See www.springbio.com for information on species reactivity predicted by sequence homology.)
DESCRIPTION:	<i>c-erbB-2</i> is a receptor tyrosine kinase of the <i>c-erbB</i> family. It is closely related in structure to the epidermal growth factor receptor. <i>c-erbB-2</i> oncoprotein is detectable in a proportion of breast and other adenocarcinomas, as well as transitional cell carcinomas.
APPLICATIONS:	Immunohistochemistry (IHC)
IHC PROCEDURE:	Specimen Preparation: Formalin-fixed, paraffin-embedded tissues are suitable for use with this primary antibody. Deparaffinization: Deparaffinize slides using xylene or xylene alternative and graded alcohols. Antibody Dilution: If using the concentrate format of this product, dilute the antibody 1:100. The dilutions are estimates; actual results may differ because of variability in methods and protocols. Antigen Retrieval: Boil tissue section in 10mM citrate buffer, pH 6.0 for 10 min followed by cooling at room temperature for 20 min. Primary Antibody Incubation: Incubate for 30 minutes at room temperature. Slide Washing: Slides must be washed in between steps. Rinse slides with PBS/0.05% Tween. Visualization: Detect the antibody as instructed by the instructions provided with the visualization system.
POSITIVE CONTROL:	Breast carcinoma
CELLULAR LOCALIZATION:	Membrane
STORAGE & STABILITY	Store at 2-8°C. Do not freeze. The user must validate any other storage conditions. When properly stored, the reagent is stable to the date indicated on the label. Do not use the reagent beyond the expiration date. There are no definitive signs to indicate instability of this product; therefore, positive and negative controls should be tested simultaneously with unknown specimens. If unexpected results are observed which cannot be explained by variations in laboratory procedures and a problem with the reagent is suspected, contact Technical Support at spring.tech@ventana.roche.com .
WARNINGS & PRECAUTIONS:	<ol style="list-style-type: none">1. Avoid contact of reagents with eyes and mucous membranes. If reagents come into contact with sensitive areas, wash with copious amounts of water.2. This product is harmful if swallowed.3. Consult local or state authorities with regard to recommended method of disposal.4. Avoid microbial contamination of reagents.